

SOLFEO, Spaceborne observations over Latin America for Emission Optimization applications (2019-2020)

European Space Agency, EO Science for Society

SOLFEO TROPOMI-based isoprene emission estimates over South America

*** Introduction**

The top-down isoprene emission estimates are derived using the adjoint of the MAGRITTE chemistry-transport model run at 0.5°x0.5° horizontal resolution (Müller et al., 2019, Bauwens et al. 2016, Stavrakou et al. 2015, Müller and Stavrakou, 2005) and constrained by tropospheric HCHO column densities from the TROPOMI satellite instrument in 2018. The HCHO data are documented in De Smedt et al. (2018) and are available via the Copernicus Open Access Hub <http://scihub.copernicus.eu>. The data were corrected for biases based on FTIR observations and comparisons with OMI HCHO data (cf. final SOLFEO report). The top-down emissions are available from <https://emissions.aeronomie.be/index.php/tropomi-based/isoprene-sa>. A short description of the project and results are summarized in the following press release: <https://eo4society.esa.int/2020/02/19/first-tropomi-based-emission-estimates-over-south-america/>.

*** File format and contents**

Monthly TROPOMI-based isoprene emissions are provided in NetCDF data format for 2018. They are expressed in kg isoprene/m²/s in a regular grid at a spatial resolution of 0.5°x0.5°. Longitudes range from 32°W to 85°W and latitudes from 15°N to 34°S.

*** Additional information**

The dataset is described in detail in the final report of the SOLFEO project.

*** References**

Bauwens, M., Stavrakou, T., Müller, J.-F., De Smedt, I., Van Roozendael, M., van der Werf,, G. R., Wiedinmyer, C., Kaiser, J., Sindelarova, K., and Guenther, A.: Nine years of global hydrocarbon emissions based on source inversion of OMI formaldehyde observations, *Atmos. Chem. Phys.*, 16, 10133-10158, <https://doi.org/10.5194/acp-16-10133-2016>, 2016.

De Smedt, I., Theys, N., Yu, H., Danckaert, T., Lerot, C., Compernolle, S., Van Roozendael, M., Richter, A., Hilboll, A., Peters, E., Pedergnana, M., Loyola, D., Beirle, S., Wagner, T., Eskes, H., van Geffen, J., Boersma, K. F., and Veefkind, P.: Algorithm theoretical baseline for formaldehyde retrievals from S5P TROPOMI and from the QA4ECV project, *Atmos. Meas. Tech.*, 11, 2395–2426, <https://doi.org/10.5194/amt-11-2395-2018>, 2018.

Müller, J.-F., T. Stavrakou, and J. Peeters: Chemistry and deposition in the Model of Atmospheric composition at Global and Regional scales using Inversion Techniques for Trace Gas Emissions (MAGRITTE v1.1). Part A. Chemical mechanism, Geosci. Model Dev., 12, 2307-2356, <https://doi.org/10.5194/gmd-12-2307-2019>, 2019.

Müller, J.-F., and T. Stavrakou: Inversion of CO and NO_x emissions using the adjoint of the IMAGES model, Atmos. Chem. Phys. 5, 1157-1186, <https://doi.org/10.5194/acp-5-1157-2005>, 2005.

Stavrakou, T., J.-F. Müller, M. Bauwens, I. De Smedt, M. Van Roozendael, M. De Mazière, C. Vigouroux, F. Hendrick, M. George, C. Clerbaux, P.-F. Coheur, and A. Guenther: How consistent are top-down hydrocarbon emissions based on formaldehyde observations from GOME-2 and OMI? Atmos. Chem. Phys., 15, 11861-11884, <https://doi.org/10.5194/acp-15-11861-2015>, 2015.

*** Contact**

Royal Belgian Institute for Space Aeronomy

Tropospheric Chemistry Modelling Group

Avenue Circulaire 3

B-1180 Brussels, BELGIUM

T. Stavrakou: jenny@aeronomie.be, tel. : +32(0)2.373.67.66

M. Bauwens: maiteb@aeronomie.be, tel. : +32(0)2.373.04.21